



Description

FOOD FOR SPECIAL MEDICAL PURPOSES

Renastep is a ready to use high energy, low volume liquid feed containing protein, carbohydrate, fat, vitamins, minerals, trace elements and docosahexaenoic acid (DHA), with low levels of potassium, phosphorus, calcium, chloride and vitamin A.

Renastep has a vanilla flavour.

Indication

Renastep is for the dietary management of kidney disease from 3 years of age onwards.

Dosage and Administration

To be determined by the clinician or dietitian and is dependent on the age, body weight and medical condition of the patient.

Preparation Guidelines

Suitable to be taken orally or via an enteral feeding tube as determined by the clinician or dietitian. Shake before use. If taken orally it is best served chilled.

Important Notice

Use under medical supervision.

Not suitable for use as a sole source of nutrition.

Suitable from 3 years of age onwards.

Regular monitoring of nutritional status and electrolyte levels is required.

For enteral use only.

Nutritional Information

		Per 100ml	Per bottle (125ml)		Per 100ml	Per bottle (125ml)	
Energy	kJ	836	1045	Minerals	Sodium	mg 84	105
		kcal	200			250	mmol
Fat	g	11.1	14	Potassium	mg	35	44
		of which saturates	g		1.7	2.1	mmol
of which monounsaturates	g	7.8	9.8	Chloride	mg	60	75
of which polyunsaturates	g	1.1	1.4		mmol	1.7	2.1
Alpha linolenic acid	mg	125	156	Calcium	mg	47	59
Docosahexaenoic acid (DHA)	mg	40	50		mmol	1.2	1.5
Linoleic acid	mg	1000	1250	Phosphorus	mg	35	44
Carbohydrate	g	21.0	26		mmol	1.1	1.4
of which sugars	g	3.4	4.3	Magnesium	mg	15	19
Protein	g	4.0	5.0		mmol	0.6	0.8
Salt	g	0.21	0.26	Trace Elements			
Vitamins				Iron	mg	2.0	2.5
Vitamin A (RE)	µg	39	49	Zinc	mg	1.2	1.5
Vitamin D	µg	1.6	2.0	Copper	mg	0.12	0.15
Vitamin E (αTE)	mg	3.6	4.5	Manganese	mg	0.20	0.25
Vitamin K	µg	7.0	8.8	Selenium	µg	5.0	6.3
Vitamin C	mg	8.0	10.0	Chromium	µg	2.5	3.1
Thiamin	mg	0.12	0.15	Molybdenum	µg	7.0	8.8
Riboflavin	mg	0.16	0.20	Iodine	µg	18	23
Niacin	mg	1.37	1.7	Others			
(Niacin Eq)	mg	(2.2)	(2.8)	L-Carnitine	mg	4.0	5.0
Vitamin B6	mg	0.21	0.26	Choline	mg	40	50
Folic acid	µg	25	31	Osmolality	600mOsm/kg		
Vitamin B12	µg	0.4	0.50				
Biotin	µg	4.0	5.0				
Pantothenic acid	mg	0.6	0.75				

Allergen Declaration

Contains **Milk** (milk protein)

Contains **Fish** (tuna oil)

Storage

Renastep should be stored in a cool dry place.

Opened: Once opened recap and refrigerate.

Discard any feed after 24 hours.

Pack Size/Weight

24 x 125ml e 3.0 L



Innovation in Nutrition
www.vitafloweb.com

A Nestlé Health Science Company
© Reg. Trademarks of Société des Produits Nestlé S.A.